

December 2001
IUT-Phys/01-11

Dirac Quantization of Some Singular Theories

A. Shirzad ¹ , P. Moyassari

*Department of Physics, Isfahan University of Technology
Isfahan, IRAN,
Institute for Studies in Theoretical Physics and Mathematics
P. O. Box: 5746, Tehran, 19395, IRAN.*

Abstract

Analyzing the constraint structure of electrodynamics, massive vector bosons, Dirac fermions and electrodynamics coupled to fermions, we show that Dirac quantization method leads to appropriate creation-annihilation algebra among the Forier coefficients of the fields.

¹e-mail: shirzad@theory.ipm.ac.ir

1 Introduction

The standard procedure for canonical quantization of fields is based on the rule

$$\{ , \}_{P.B} \rightarrow \frac{1}{i\hbar} [,] \quad (1)$$

For simple examples such as Klien-Gordon field, the above rule leads to creation-annihilation algebra between coefficients of Forier modes of the fields. However, the problem is not so simple in most cases of interest such as gauge theories. In fact a large number of models come from singular Lagrangians, where a well-defined Poisson bracket is far from reach. In these cases, the traditional method for quantization is as follows: One finds first the *physical fields* in some sense and considers their Forier transformations; and then *assumes* that coefficients of Forier modes act as creation-annihilation operators [7, 8]. Usually, there exist difficulties in recognizing the physical fields such that a standard method is not followed by authors to find them. On the other hand, no justifiable reason is given for assuming the creation-annihilation algebra among coefficients of Forier expansion. Accordingly we think that this approach violates the unified quantization method given by (1). However, it is well-known that a well-defined bracket, known as Dirac bracket, can be assigned to singular Lagrangians. In this method the quantization rule (1) is generalized to

$$\{ , \}_{D.B} \longrightarrow \frac{1}{i\hbar} [,]. \quad (2)$$

This rule reduces simply to (1) for regular models.

As far as we know, this unified method for quantization, i.e. Dirac quantization, is not exactly followed in the literature for models of physicists' interest.

Our main objective in this paper is that the appropriate creation-annihilation algebra would emerge directly by following the Dirac method, with no need to any ad-hoc assumption. After a brief review of Dirac quantization in the next section, we will do this job for electrodynamics, massive vector bosons, Dirac fermion and electrodynamics coupled to fermions in sections 3-6.

2 Dirac quantization

The Lagrangian $L(q, \dot{q})$ is said to be singular if the momenta

$$p_i = \frac{\partial L}{\partial \dot{q}_i} \quad (3)$$

are not independent functions of $(q, \dot{q})$. It follows that a number of functions $\phi_i(p, q)$ should vanish as primary constraints. It can be shown [1] that time development of any function $g(q, p)$ is achieved via the equation

$$\dot{g} = \{g, H_T\} \quad (4)$$

where

$$H_T = H_C + \lambda^i \phi_i \quad (5)$$

in which H_C is the canonical Hamiltonian and λ^i are Lagrange multipliers. Primary constraints should vanish at all times. So the consistency conditions

$$\dot{\phi}_i = \{\phi_i, H_C\} + \lambda^j \{\phi_i, \phi_j\} \approx 0 \quad (6)$$

should be imposed on them, where $\approx$ means weak equality, i.e. equality on the constraint surface. The consistency conditions (6) may lead to determining a number of Lagrange multipliers and/or emerging some new constraints, known as secondary constraints. The procedure should be repeated until the last step where the maximum number of Lagrange multipliers are determined and the consistency of all remaining constraints are guaranteed on the constraint surface.

The whole system of constraints divide to first class constraints ϕ_a and second class constraints χ_n such that

$$\begin{aligned} \{\phi_a, \phi_{a'}\} &\approx 0 \\ \{\phi_a, \chi_n\} &\approx 0 \\ \{\chi_n, \chi_m\} &\approx c_{nm} \quad \det(c_{nm} \neq 0). \end{aligned} \quad (7)$$

The second class constraints, which are even in number, reduce the dimensionality of phase space and lead to determining a number of the Lagrange multipliers [1]. While, the first class constraints are generators of gauge

transformations [2]. In order to fix the gauges one should impose a number of gauge fixing conditions $\Omega_{a'}(q, p) = 0$ on the system. For gauges to be fixed completely, the determinant of $\{\Omega_{a'}, \phi_a\}$ should not vanish [3]. In other words, after fixing the gauges the set of constraints and gauge fixing conditions, i.e.

$$\Phi_A \equiv (\chi_n; \phi_a, \Omega_{a'}), \quad (8)$$

should act as a system of second class constraints. The reduced phase space $\Phi_A \approx 0$ possesses a well-defined bracket, known as Dirac bracket, which is defined for any two arbitrary functions $f(q, p)$ and $g(q, p)$ as

$$\{f, g\}_{D.B} = \{f, g\}_{P.B} - \{f, \Phi_A\}_{P.B} C^{AB} \{\Phi_B, g\}_{P.B} \quad (9)$$

where C^{AB} is the inverse of $C_{AB} = \{\Phi_A, \Phi_B\}_{P.B}$. In this way all redundant coordinates are omitted and just the physical degrees of freedom are remained in the reduced phase space. Then one can quantize the system in a consistent way by transforming the Dirac brackets into commutators, as mentioned before in (2). Let us now follow the procedure for some important physical models.

3 Electrodynamics

The Lagrangian density for electrodynamics reads

$$\mathcal{L} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} \quad (10)$$

where

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (11)$$

From definition of momentum fields, i.e.

$$\Pi^\mu(x) = \frac{\delta L}{\delta(\partial_0 A_\mu(x))} = -F^{0\mu}(x), \quad (12)$$

one finds that $\Pi^0(x)$ vanishes. So the primary constraint of the system is

$$\phi^1(x) = \Pi^0(x). \quad (13)$$

To quantize the theory most authors replace the main Lagrangian with Fermi Lagrangian:

$$\mathcal{L}_f = -\frac{1}{2}(\partial_\mu A_\nu)(\partial^\mu A^\nu) \quad (14)$$

which, together with the Loran condition $\partial_\mu A^\mu = 0$, gives the same equations of motion [7, 9]. It is well-known that the Fermi Lagrangian leads to negative norm states after quantization, first problem; and the Loran gauge condition can not be imposed in the operator form, next problem. To this end, one can show that imposing the condition

$$\langle \text{phy} | \partial_\mu A^\mu | \text{phy} \rangle = 0 \quad (15)$$

on physical states solves both problems. However, it seems that the above method is adjusted just for the problem of quantization of electrodynamics. While it is preferable to follow a general procedure to quantize all models.

Now we show that following the Dirac method, one can quantize the theory in a straightforward method. Returning to the main Lagrangian(10), the canonical Hamiltonian reads

$$H_C = \int \left(\frac{1}{2} \Pi_i \Pi^i + A_0 \partial_i \Pi^i + \frac{1}{4} F_{ij} F^{ij} \right) d^3x \quad (16)$$

Consistency of ϕ^1 from (6) gives the second level constraint

$$\phi^2 \equiv \{ \phi^1, H_C \} = \partial_i \Pi^i \quad (17)$$

It is easy to see that $\{ \phi^2, \phi^1 \} \approx 0$ and $\{ \phi^2, H_C \} \approx 0$. Therefore, there is only one constraint chain as

$$\begin{aligned} \phi^1 &= \Pi^0 \\ \phi^2 &= \partial_i \Pi^i \end{aligned} \quad (18)$$

As discussed in [3], to fix the gauge it is enough to impose the gauge fixing condition

$$\Omega^2 = \partial_i A^i. \quad (19)$$

Since $\{ \Omega^2, \phi^2 \} \neq 0$, Ω^2 fixes the gauge transformation generated by ϕ^2 . Consistency of gauge fixing condition Ω^2 gives the other gauge fixing condition conjugate to ϕ^1 :

$$\Omega^1 \equiv \{ \Omega^2, H_T \} \approx \partial^i \partial_i A_0 \quad (20)$$

Finally, one should calculate the Dirac brackets due to system of constraints

$$(\phi^1, \phi^2; \Omega^2, \Omega^1). \quad (21)$$

After a little algebra, the Dirac brackets emerges as

$$\begin{aligned}
[A^\mu(\mathbf{x}, t), \Pi_\nu(\mathbf{x}', t)]_{D.B} &= (\delta^\mu{}_\nu - g_{\nu 0} \delta^\mu{}_0) \delta(\mathbf{x} - \mathbf{x}') - \partial^\mu \partial_\nu \frac{1}{4\pi|\mathbf{x} - \mathbf{x}'|} \\
[A^\mu(\mathbf{x}, t), A_\nu(\mathbf{x}', t)]_{D.B} &= 0 \\
[\Pi^\mu(\mathbf{x}, t), \Pi_\nu(\mathbf{x}', t)]_{D.B} &= 0
\end{aligned} \tag{22}$$

Assuming the system to be contained in a box of volume V , the most general solution to equations of motion reads

$$A^\mu(x) = \sum_k \sum_{r=0}^3 \frac{1}{(2Vw_k)^{1/2}} (\epsilon_r^\mu(\mathbf{k}) a_r(\mathbf{k}) e^{-ikx} + \epsilon_r^\mu(\mathbf{k}) a_r^\dagger(\mathbf{k}) e^{ikx}) \tag{23}$$

Where $w_k = |\mathbf{k}|$ and the polarization vectors $\epsilon_r^\mu(\mathbf{k})$ can be chosen to be

$$\begin{aligned}
\text{scalar :} \quad & \epsilon_0^\mu(\mathbf{k}) = (1, 0, 0, 0) \\
\text{transverse :} \quad & \epsilon_{1,2}^\mu(\mathbf{k}) = (0, \mathbf{e}(\mathbf{k})) \quad \mathbf{k} \cdot \mathbf{e}(\mathbf{k}) = 0 \\
\text{longitudinal :} \quad & \epsilon_3^\mu(\mathbf{k}) = (0, \frac{\mathbf{k}}{|\mathbf{k}|})
\end{aligned} \tag{24}$$

Using (12), the momentum fields can also be written as

$$\Pi^\mu(x) = \sum_k \sum_{r=0}^3 \frac{i}{(2Vw_k)^{1/2}} (k^\mu \epsilon_r^0(\mathbf{k}) - w_k \epsilon_r^\mu(\mathbf{k})) (-a_r(\mathbf{k}) e^{-ikx} + a_r^\dagger(\mathbf{k}) e^{ikx}) \tag{25}$$

Using (23) and (25), one can find $a_r(\mathbf{k})$ in term of fields (at $t = 0$) as

$$\xi_r a_r(\mathbf{k}) = \frac{1}{(2w_k)^{1/2}} \int d^3x \epsilon_{\mu r} (-i\Pi^\mu - w_k A^\mu) e^{i\mathbf{k} \cdot \mathbf{x}} + \frac{k^j \epsilon_{jr}}{w_k} a_0(\mathbf{k}) \tag{26}$$

where $\xi_0 = -1$ and $\xi_{1,2,3} = 1$. Finally, from (22), (26) and its complex conjugate for $a_r^\dagger(\mathbf{k})$ one can derive the Dirac brackets among $a_r(\mathbf{k})$ and $a_r^\dagger(\mathbf{k})$ as:

$$\begin{aligned}
[a_0(\mathbf{k}), a_0^\dagger(\mathbf{k}')]_{D.B} &= 0 \\
[a_3(\mathbf{k}), a_3^\dagger(\mathbf{k}')]_{D.B} &= 0 \\
[a_r(\mathbf{k}), a_s^\dagger(\mathbf{k}')]_{D.B} &= -i\delta_{rs}\delta_{\mathbf{k},\mathbf{k}'} \quad r, s = 1, 2
\end{aligned} \tag{27}$$

Now everything is ready for quantization. It is just needed to use quantization rule (2) to find commutators. As is apparent the scalar and longitudinal polarizations disappear in a consistent way, and $a_{1,2}(\mathbf{k})$ and $a_{1,2}^\dagger(\mathbf{k})$ obey creation-annihilation algebra. In other words, just transverse photons remain as physical degrees of freedom in the quantized theory.

4 Massive Vector Bosons

The model is given by the Lagrangian density

$$\mathcal{L} = -\frac{1}{2}F_{\mu\nu}^\dagger F^{\mu\nu} + m^2 W^{\dagger\mu} W_\mu \quad (28)$$

where the complex fields W^μ and $W^{\mu\dagger}$ describe vector bosons of mass m and

$$F_{\mu\nu} = \partial_\mu W_\nu - \partial_\nu W_\mu. \quad (29)$$

The equations of motion reads

$$\partial_\mu F^{\mu\nu} + m^2 W^\nu = 0. \quad (30)$$

(similar relations as (29) and (30) can be understood for $W^{\dagger\mu}$ and $F^{\dagger\mu\nu}$). Taking the divergence of both sides of (30) gives

$$\partial_\nu W^\nu = 0 \quad (31)$$

In the context of the theory of constrained systems $\partial_\nu W^\nu$ (as well as $\partial_\nu W^{\dagger\nu}$) is recognized as a Lagrangian constraint [6]. It shows that the fields W^μ are not independent. Traditionally, people consider the scalar polarization as the redundant field and quantize the remaining ones (i.e. transverse and Longitudinal polarizations) as physical degrees of freedom. But usually there is no justification that "why scalar polarization should be omitted?". Now, using the theory of constrained systems, we show that this is really the case. From (28) the canonical momentum fields are $\Pi^\mu = F^{\dagger\mu 0}$ and $\Pi^{\dagger\mu} = F^{\mu 0}$. This leads to primary constraints

$$\phi_1^1 = \Pi^0 \quad \phi_2^1 = \Pi^{\dagger 0} \quad (32)$$

The canonical Hamiltonian reads

$$H_C = \int d^3x \left(\Pi_i^\dagger \Pi^i + \partial_i \Pi^i W_0 + \partial_i \Pi^{\dagger i} W_0^\dagger + \frac{1}{2} F^{ij} F_{ij}^\dagger - m^2 W_\mu^\dagger W^\mu \right) \quad (33)$$

Consistency conditions (6) on primary constraints gives the second level constraints as

$$\phi_1^2 = \partial_i \Pi^i + m^2 W_0^\dagger \quad \phi_2^2 = \partial_i \Pi^{\dagger i} + m^2 W_0 \quad (34)$$

It is easy to see that we have four second class constraints. In fact since $\{\phi_1^2, \phi_2^1\} \neq 0$ and $\{\phi_2^2, \phi_1^1\} \neq 0$ we have two *cross conjugate* second class chains, in the terminology of [5]. The Dirac brackets can be derived straightforwardly as

$$\begin{aligned} [W_i(\mathbf{x}, t), \Pi^j(\mathbf{x}', t)]_{D.B} &= [W_i^\dagger(\mathbf{x}, t), \Pi^{\dagger j}(\mathbf{x}', t)]_{D.B} = \delta_i^j \delta(\mathbf{x} - \mathbf{x}') \\ [W_0(\mathbf{x}, t), \Pi^0(\mathbf{x}', t)]_{D.B} &= [W_0^\dagger(\mathbf{x}, t), \Pi^{\dagger 0}(\mathbf{x}', t)]_{D.B} = 0 \end{aligned} \quad (35)$$

As in electrodynamics, the solution of equations of motion reads

$$W^\mu(x) = \sum_k \sum_{r=0}^3 \frac{1}{(2V w_k)^{1/2}} [\epsilon_r^\mu(\mathbf{k}) a_r(\mathbf{k}) e^{-ikx} + \epsilon_r^\mu(\mathbf{k}) b_r^\dagger(\mathbf{k}) e^{ikx}] \quad (36)$$

where $w_k = \sqrt{k^2 + m^2}$. Using (36), its complex conjugate for $W_\mu^\dagger$, and expression for Π_μ and $\Pi_\mu^\dagger$, one finds

$$\begin{aligned} \xi_r a_r &= \frac{1}{(2w_k)^{1/2}} \int d^3x \epsilon_{\mu r}(\mathbf{k}) (-i\Pi^{\dagger \mu}(\mathbf{x}) - w_k W^\mu(\mathbf{x})) e^{i\mathbf{k} \cdot \mathbf{x}} + \frac{k^j \epsilon_{jr}(\mathbf{k})}{w_k} a_0(\mathbf{k}) \\ \xi_r b_r &= \frac{1}{(2w_k)^{1/2}} \int d^3x \epsilon_{\mu r}(\mathbf{k}) (-i\Pi^\mu(\mathbf{x}) - w_k W^{\dagger \mu}(\mathbf{x})) e^{i\mathbf{k} \cdot \mathbf{x}} + \frac{k^j \epsilon_{jr}(\mathbf{k})}{w_k} b_0(\mathbf{k}) \end{aligned} \quad (37)$$

where the fields are considered at $t = 0$. The expression for $a_r^\dagger(\mathbf{k})$ and $b_r^\dagger(\mathbf{k})$ can be written by complex conjugating the above formulas. Then using the Dirac brackets (35) the following fundamental Dirac brackets emerge

$$\begin{aligned} [a_r(\mathbf{k}), a_s^\dagger(\mathbf{k}')]_{D.B} &= -i\delta_{rs}\delta_{kk'} \quad r, s = 1, 2, 3 \\ [b_r(\mathbf{k}), b_s^\dagger(\mathbf{k}')]_{D.B} &= -i\delta_{rs}\delta_{kk'} \quad r, s = 1, 2, 3 \\ [a_0(\mathbf{k}), a_0^\dagger(\mathbf{k}')]_{D.B} &= 0 \\ [b_0(\mathbf{k}), b_0^\dagger(\mathbf{k}')]_{D.B} &= 0 \end{aligned} \quad (38)$$

As is apparent, after quantization according to (2), the scalar polarization disappears. Moreover $a_r(\mathbf{k})$ and $a_r^\dagger(\mathbf{k})$ act as annihilation and creation operators for W^+ particle with transverse and longitudinal polarizations, while $b_r(\mathbf{k})$ and $b_r^\dagger(\mathbf{k})$ act in the same way for W^- particle.

5 Dirac Fermions

Fermion fields satisfy different algebra, compared to bosons, even at the classical level. In fact the Poisson bracket of two functions on a super phase space have generalized algebraic properties [10], among them we remind the following:

$$\begin{aligned}\{F, G\} &= (-1)^{\epsilon_F \epsilon_G + 1} \{G, F\} \\ \{F, GH\} &= \{F, G\} H + (-1)^{\epsilon_F \epsilon_G} G \{F, H\}\end{aligned}\tag{39}$$

where ϵ_F and ϵ_G are Grassmanian parity of F and G respectively. This shows specially that the Poisson bracket of two fermionic variables is even under changing the order. As we will show, this leads to defining anticommutators, instead of commutators, for the operators at the quantum level. The Dirac fields are introduced within the Lagrangian density

$$\mathcal{L} = \bar{\Psi}(x)(i\gamma^\mu \partial_\mu - m)\Psi(x)\tag{40}$$

where $\Psi(x)$ and $\bar{\Psi}(x)$ are fermionic fields. The canonical momentum fields are derived as

$$\begin{aligned}\Pi(x) &= \frac{\partial^L L}{\partial \dot{\Psi}} = -i\bar{\Psi}\gamma^0 \\ \bar{\Pi}(x) &= \frac{\partial^L L}{\partial \dot{\bar{\Psi}}} = 0\end{aligned}\tag{41}$$

where ∂^L means left differentiation. From (41) the following primary constraints emerge

$$\phi_1 = \Pi + i\bar{\Psi}\gamma^0 \quad \phi_2 = \bar{\Pi}\tag{42}$$

As is obvious, $\{\phi_1, \phi_2\} \neq 0$, showing that ϕ_1 and ϕ_2 are second class. Therefore, the consistency condition on them just determines the Lagrang multipliers. Using the fundamental Poisson brackets of fermion fields, i.e.

$$\{\Psi(\mathbf{x}, t), \Pi(\mathbf{x}', t)\} = \{\bar{\Psi}(\mathbf{x}, t), \bar{\Pi}(\mathbf{x}', t)\} = -\delta(\mathbf{x} - \mathbf{x}'),\tag{43}$$

the Dirac berackets, with the same algebraic properties as Poisson bracket, are simply derived as

$$\begin{aligned}
[\Psi(\mathbf{x}, t), \bar{\Psi}(\mathbf{x}', t)]_{D.B} &= -i\gamma^0 \delta^3(x - x') \\
[\Psi(\mathbf{x}, t), \Pi(\mathbf{x}', t)]_{D.B} &= -\delta^3(x - x') \\
[\bar{\Psi}(\mathbf{x}, t), \bar{\Pi}(\mathbf{x}', t)]_{D.B} &= 0 \\
[\Pi(\mathbf{x}, t), \bar{\Pi}(\mathbf{x}', t)]_{D.B} &= 0
\end{aligned} \tag{44}$$

The famous Forier expansion of Dirac fields, read

$$\begin{aligned}
\Psi(x) &= \sum_k \sum_{r=1}^2 \left(\frac{m}{VE_p}\right)^{1/2} \left[c_r(\mathbf{p}) u_r(\mathbf{p}) e^{-ipx} + d_r^\dagger(\mathbf{p}) v_r(\mathbf{p}) e^{ipx} \right] \\
\bar{\Psi}(x) &= \sum_k \sum_{r=1}^2 \left(\frac{m}{VE_p}\right)^{1/2} \left[c_r^\dagger(\mathbf{p}) \bar{u}_r(\mathbf{p}) e^{ipx} + d_r(\mathbf{p}) \bar{v}_r(\mathbf{p}) e^{-ipx} \right]
\end{aligned} \tag{45}$$

where $u_r(\mathbf{p})$ and $v_r(\mathbf{p})$ are Dirac spinors. At this point text books on quantum field theory usually *assume* the following anti-commutator algebra between Forier coefficients as the beginning point for quantization of Dirac fermions

$$\begin{aligned}
[c_r(\mathbf{p}), c_s^\dagger(\mathbf{p}')]_+ &= \delta_{rs} \delta_{pp'} \\
[d_r(\mathbf{p}), d_s^\dagger(\mathbf{p}')]_+ &= \delta_{rs} \delta_{pp'}
\end{aligned} \tag{46}$$

However, using

$$\begin{aligned}
2\sqrt{m/E_p} c_r(\mathbf{p}) &= \int d^3x \bar{u}_r(\mathbf{p}) \Psi(\mathbf{x}, 0) e^{i\mathbf{p}\cdot\mathbf{x}} \\
-2\sqrt{m/E_p} d_r(\mathbf{p}) &= \int d^3x \bar{\Psi}(\mathbf{x}, 0) v_r(\mathbf{p}) e^{i\mathbf{p}\cdot\mathbf{x}}
\end{aligned} \tag{47}$$

and (44) it is easy to see that

$$\begin{aligned}
[c_r(\mathbf{p}), c_s^\dagger(\mathbf{p}')]_{D.B} &= -i\delta_{rs} \delta_{pp'} \\
[d_r(\mathbf{p}), d_s^\dagger(\mathbf{p}')]_{D.B} &= -i\delta_{rs} \delta_{pp'}
\end{aligned} \tag{48}$$

Note that the above (classical) berackets are even under exchange of variables. It is reasonable to keep the algebraic properties of classical berackets

at quantum level. So in the quantization rule (2) we should replace anti-commutators on the right hand side, in case of fermions. The final result is the creation -annihilation algebra(46). We see once again that Dirac method for quantization gives the correct algebra of quantum operators with no need to unjustified assumptions.

6 Electrodynamics coupled to Fermions

The Lagrangian density for electromagnetic field coupled to Dirac fermions is known as

$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \bar{\Psi}(i\gamma^\mu\partial_\mu - m)\Psi - e\bar{\Psi}\gamma^\mu A_\mu\Psi \quad (49)$$

The system of constraints can be obtained similar to the case of free electrodynamics and Dirac fermions as follows:

$$\begin{aligned} \phi_1^1 &= \Pi^0 & \phi_2 &= \bar{\Pi} & \phi_3 &= \Pi + i\bar{\Psi}\gamma^0\Psi \\ \phi_1^2 &= -\partial_i\Pi^i + ie(\bar{\Psi}\bar{\Pi} + \Pi\Psi) \end{aligned} \quad (50)$$

where $\phi_1^2 = \{\phi_1^1, H_T\}$. As before, ϕ_2 and ϕ_3 are second class; however, ϕ_1^1 and ϕ_1^2 are elements of a first class chain. Parenthetically, the infinitesimal gauge transformation

$$\begin{aligned} A_\mu &\longrightarrow A_\mu + \partial_\mu\zeta \\ \Psi &\longrightarrow (1 - ie\zeta)\Psi \\ \bar{\Psi} &\longrightarrow (1 + ie\zeta)\bar{\Psi}, \end{aligned} \quad (51)$$

can be obtained as the action of the gauge generator [2, 4]

$$G = \int d^3x(\dot{\zeta}\phi_1^1 - \zeta\phi_1^2). \quad (52)$$

In order to fix the gauge, following the method of [3], we can introduce the gauge fixing condition

$$\Omega_1^2 = \partial_i A^i \quad (53)$$

conjugate to the last element ϕ_1^2 . Then consistency of Ω_1^2 gives the next gauge fixing condition as

$$\Omega_1^1 = \partial_i\partial^i A_0 \quad (54)$$

The whole system of constraints $(\phi_1^1, \phi_1^2, \Omega_1^2, \Omega_1^1; \phi_2, \phi_3)$ imply the following Dirac brackets

$$\begin{aligned}
[A^\mu(\mathbf{x}, t), \Pi_\nu(\mathbf{x}', t)]_{D.B} &= (\delta^\mu{}_\nu - g_{\nu 0} \delta^\mu{}_0) \delta(\mathbf{x} - \mathbf{x}') - \partial^\mu \partial_\nu \frac{1}{4\pi|\mathbf{x} - \mathbf{x}'|} \\
[\Psi(\mathbf{x}, t), \bar{\Psi}(\mathbf{x}', t)]_{D.B} &= -i\gamma^0 \delta(\mathbf{x} - \mathbf{x}') \\
[\Psi(\mathbf{x}, t), \Pi(\mathbf{x}', t)]_{D.B} &= -\delta(\mathbf{x} - \mathbf{x}').
\end{aligned} \tag{55}$$

All other Dirac brackets vanish. These are the same brackets as obtained for free electrodynamics and free Dirac fermions in (22) and (44). This shows that the algebra of quantum fields are the same as before. But some care is necessary to obtain the appropriate algebra among the coefficients of Forier expansion. That is because in the presence of coupling term $-e\bar{\Psi}\gamma^\mu A_\mu\Psi$ the equations of motion are different from that for free fields and the Forier expansions (23) and (45) are no longer valid. However, in quantum theory the coupling term $-e\bar{\Psi}\gamma^\mu A_\mu\Psi$ is considered as a perturbation. So the quantum fields obey the same equations of motion as obtained from the free part of the Lagrangian (i.e. the first two terms in (49)). Consequently the fields A^μ , Ψ and $\bar{\Psi}$ have the same Forier expansions as in the free cases. To this end, beginning with quantum versions of (55) one is allowed to use free field expansion (23) and (45) to obtain the same creation-annihilation algebra as in (27) and (48).

References

- [1] Dirac P A M, "*Lectures notes on Quantum Mechanics*" New York: Yeshiva University Press, 1964
- [2] Gracia X and Pons J M, *Ann.Phy* **187** (1988) 355
- [3] Shirzad A and Loran F, *Hep-th* 9912289
- [4] Shirzad A and Shabani Moghaddam M, *J. Phys* **A32** (1999) 8185
- [5] Shirzad A and Loran F, *Hep-th* 0003010
- [6] Shirzad A, *J.Phys* **A31**(1998) 2747
- [7] Mandl F and Show G, "*Quantum Field Theory*" Manchester University, 1988

- [8] Ryder L, "*Quantum Field Theory*" Cambridge University Press, 1985
- [9] Zuber J B and Itzikson C, "*Quantum Field Theory*" The United states of America ,1980
- [10] Henneaux M and Teitelboim C, "*Quantization of Gauge System*" Princeton University Press, 1992